



Chemical Compositions, and Antimicrobial and Mosquito Larvicidal Activities of Essential Oils from Four *Syzygium* Species *Syzygium formosum* (Wall.) Masam., *S. syzygioides* (Miq.) Merr. & L.M. Perry, *S. megacarpum* (Craib) Rathakr. & N.C. Nair, and *S. chantaranothaianum* W.K. Soh & J. Parn

Le Thi Huong, Hoang Vinh Phu, Nguyen Huy Hung, Le Duc Giang, Do Ngoc Dai, Nguyen Quang Hop & Ninh The Son

To cite this article: Le Thi Huong, Hoang Vinh Phu, Nguyen Huy Hung, Le Duc Giang, Do Ngoc Dai, Nguyen Quang Hop & Ninh The Son (10 Mar 2024): Chemical Compositions, and Antimicrobial and Mosquito Larvicidal Activities of Essential Oils from Four *Syzygium* Species *Syzygium formosum* (Wall.) Masam., *S. syzygioides* (Miq.) Merr. & L.M. Perry, *S. megacarpum* (Craib) Rathakr. & N.C. Nair, and *S. chantaranothaianum* W.K. Soh & J. Parn, Journal of Essential Oil Research, DOI: [10.1080/10412905.2024.2322593](https://doi.org/10.1080/10412905.2024.2322593)

To link to this article: <https://doi.org/10.1080/10412905.2024.2322593>



Published online: 10 Mar 2024.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)



Chemical Compositions, and Antimicrobial and Mosquito Larvicidal Activities of Essential Oils from Four *Syzygium* Species *Syzygium formosum* (Wall.) Masam., *S. syzygioides* (Miq.) Merr. & L.M. Perry, *S. megacarpum* (Craib) Rathakr. & N.C. Nair, and *S. chantaranthaianum* W.K. Soh & J. Parn

Le Thi Huong^a, Hoang Vinh Phu^a, Nguyen Huy Hung^{b,c}, Le Duc Giang^{ib,d}, Do Ngoc Dai^e, Nguyen Quang Hop^f and Ninh The Son^{g,h}

^aFaculty of Biology, College of Education, Vinh University, Vinh, Vietnam; ^bCenter for Advanced Chemistry, Institute of Research and Development, Duy Tan University, Da Nang, Vietnam; ^cDepartment of Pharmacy, Duy Tan University, Da Nang, Vietnam; ^dFaculty of Chemistry, College of Education, Vinh University, Vinh, Vietnam; ^eFaculty of Agriculture, Forestry and Fishery, Nghe An University of Economics, Vinh, Vietnam; ^fFaculty of Chemistry, Hanoi Pedagogical University 2 (HPU2), Nguyen Van Linh, Vietnam; ^gInstitute of Chemistry, Vietnam Academy of Science and Technology (VAST), Hanoi, Vietnam; ^hDepartment of Chemistry, Graduate University of Science and Technology, VAST, Hanoi, Vietnam

ABSTRACT

Chemical compositions in the leaf essential oils of four *Syzygium* species were first identified by the GC-FID/MS analysis. Monoterpene hydrocarbons (39.6–51.1%), and sesquiterpene hydrocarbons (27.0–37.8%) were the main chemical classes in *S. formosum* and *S. syzygioides* leaf oils, whereas sesquiterpene hydrocarbons (83.0–83.8%) were predominant in *S. Megacarpum* and *S. chantaranthaianum* leaf essential oils. Bicyclogermacrene (9.1–37.0%) was the principal compounds in these essential oils. All the tested essential oils with the MIC values of 16–128 µg/mL were comparable to the positive control streptomycin against Gram-positive bacteria *Staphylococcus aureus* and *Bacillus cereus*, and Gram-negative bacterium *Pseudomonas aeruginosa*. The leaf essential oils of *S. formosum*, *S. syzygioides*, and *S. chantaranthaianum* with the MIC values of 16–128 µg/mL were comparable to the positive control cycloheximide against the yeast *Candida albicans*. Four samples also exhibited good larvicidal activity against the mosquito *Aedes aegypti* with the 24-h and 28-h LC₅₀ values of 25–40 µg/mL

ARTICLE HISTORY

Received 17 November 2022
Accepted 16 February 2024

KEYWORDS

Syzygium; essential oils; antimicrobial; mosquito larvicidal

1. Introduction

Numerous factors can influence the synthesis, yield, and composition of chemical components in essential oils, as well as their existence from inside the plants until their final isolation. By mean of this, the influential aspects have been researched, especially for economically significant crops, in an effort to maximize the growing conditions, and harvest timing while achieving increased yields of premium essential oils that meet market demands (1–3). Therefore, it is crucial to understand the variables that affect species' chemical variability and production. Physiological changes, environmental influences, geographic variances, genetic factors, plant material/space requirements, and the necessity for time collection are a few of these (4,5).

Syzygium is one of the largest genera in the family Myrtaceae with ca. 1200 evergreen trees and shrubs (6). The plants of this genus were recorded to distribute from Africa through Asia, Malesia and Australia to the Pacific islands (6). The Clove (*S. aromaticum*) is a well-known economically crucial species, in which its unopened

flower buds are used as a precious spice (7). Medicinally, *Syzygium* plant extracts and isolated compounds have also a broad panel of pharmacological activities, such as antimicrobial, antidiabetic, anti-inflammatory, and nephroprotective activities (7,8). It is also recognized that *Syzygium* plants are rich in essential oils containing terpenoid compounds. As an example, the antibacterial essential oil of *S. cumini* leaves was characterized by the presence of major components pinocarveol (15.1%), and α -terpeneol (8.9%) (9). Among 49 species recorded in Vietnam, some of them have been objects in phytochemical studies to identify chemical profiles in their essential oils. β -Caryophyllene (42.53–64.53%) can be seen as the main compound in the leaf essential oils of two Vietnamese *Syzygium* species *S. caryophyllatum* and *S. lineatum*, whereas the leaf essential oil of *S. hancei* was represented by γ -guaiene (11.07%) (10).

Regarding biological activities, *Syzygium* species showed potentials in antimicrobial and mosquito larvicidal treatments. The leaf essential oil of *S. grande*

inhibited the growth of *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus vulgaris*, *P. aeruginosa*, *S. aureus*, and *B. subtilis* with the MIC values of 0.25–0.75 mg/mL (11). The Clove essential oil controlled wound infections in mice caused by methicillin resistant *S. aureus* via its interaction with imipenem (12). *S. zeylanicum* leaf essential oil exhibited remarkably effective and eco-friendly larvicides against *Anopheles subpictus*, *An. albopictus*, and *Culex tritaeniorhynchus* larvae (13). The Clove essential oil was also reported to protect against *An. stephensi* larvae with the LC₅₀ and LC₉₀ values of 57.49 and 93.14 ppm, respectively (14).

The current study aims to report the chemical identification of essential oils of four wild *Syzygium* species, collected from the North Central Coast region of Vietnam, including *S. formosum* (Wall.) Masam. (local name: Trâm lá chùm ba), *Syzygium syzygioides* (Miq.) Merr. & Perry (Trâm kiền kiền), *S. megacarpum* (Craib) Rathakr. & N.C.Nair (Trâm lá lớn), and *Syzygium chantaranothaianum* W. K. Soh & J. Parn. (Trâm chan lốt), as well as their antimicrobial, and mosquito larvicidal activities.

2. Materials and methods

2.1. Plant materials

The fresh leaves of four studied species (7-year-old plants) were collected from Pu Hoat Natural Reserve, Nghean, Vietnam in 04/2022. The Latin names were identified by the co-author Le Thi Huong. The geographic coordinates included *S. formosum* (19°40'3"N and 104°55'30"), *S. syzygioides* (19°44'36" and 104°48'2"), *S. megacarpum* (19°48'31" and 105°5'47"), and *S. chantaranothaianum* (19°42'21" and 104°50'2"). The voucher specimens, including SL-01 (*S. formosum* leaves), SL-02 (*S. syzygioides* leaves), SL-03 (*S. megacarpum* leaves), and SL-04 (*S. chantaranothaianum* leaves), have been deposited in Faculty of Biology, College of Education, Vinh University. The obtained samples (2.5 kg each) were immediately cut into pieces, and immersed in distilled water at a ratio of 1:1, w/v. They were then hydro-distilled using a Clevenger apparatus for 3.5 h to give the yellow essential oils. The yields of extraction, which were calculated following dried materials, reached a range of 0.15–0.25%.

2.2. GC-FID/MS analysis

Gas chromatography with flame ionization detection (GC-FID) was carried out following the conditions (15–18): Agilent Technologies HP-5 MS column (30 m x 0.25 mm, film thickness 0.25 µm), Helium carrier gas (1.1 mL/min), injector temperature of 260°C, detector

temperature of 270°C, column temperature program: 65°C (3 min hold), increase to 230°C (4°C/min), 230°C (10 min hold), inlet pressure of 6.0 kPa, split mode injection (split ratio, 10:1), 1.1 µL injection volume.

Gas chromatography-mass spectrometry (GC-MS) was performed in the same manner: Agilent Technologies HP 7890A Plus Chromatograph (Santa Clara, CA, USA), HP-5 MS (30 m x 0.25 mm, film thickness 0.25 µm) column, HP 5973 MSD mass detector, Helium carrier gas (1.1 mL/min), MS ionization voltage of 70 eV, emission current of 40 mA, acquisitions range of 40–400 amu, a sampling rate of 1.0 scan/s. The GC was operated under the same circumstances as GC-FID. The retention indices (RI) based on a series of n-alkanes, co-injection with pure compounds (Sigma-Aldrich, St. Louis, MO, USA) or identified essential oil components, MS search (NIST 17 and Wiley 10th Version libraries), and comparison with the literature MS fragmentation were used to identify the chemical components of the essential oils (15–18). It was mainly based on the GC peak area (FID response) and without the use of correction factors, the relative concentrations (%) of the constituents were computed. The measurements were made three times.

2.3. Antimicrobial assay

Antimicrobial effect of six leaf essential oils was performed using the broth dilution method (19). Seven pathogenic bacterial strains have been used, including three Gram-positive bacterial strains *Enterococcus faecalis* ATCC299212, *Staphylococcus aureus* ATCC25923, and *Bacillus cereus* ATCC14579, three strains of Gram-negative bacterial strains *Escherichia coli* ATCC 25,922, *Pseudomonas aeruginosa* ATCC27853, and *Salmonella enterica* ATCC13076, and one yeast strain *Candida albicans* ATCC 10,231.

The selection of investigated concentrations was based on our previous publication (19), in which the tested essential oil was active with the specific concentration ranges. Stock solution of essential oil was prepared by DMSO (1%). Dilution series (2-fold) were prepared from 512 to 16 µg/mL by distilled water. They were then transferred to 96-well plates. Bacteria grown in double-strength Mueller-Hinton broth were standardized to 5×10^5 CFU/mL. The last row of well plates containing only the serial dilutions of samples without microorganisms was used as a positive control (no growth). Distilled water and medium served as a negative control (no antimicrobial agent). Streptomycin and nystatin were used as standards for antibacterial and anti-yeast activities, respectively. Experiments were repeated in triplicate. The results

were displayed by the MIC values (the lowest dose at which bacterial growth is totally inhibited).

2.4. Mosquito larvicidal assay

Eggs of *Ae. aegypti* were purchased from Institute of Biotechnology, VAST, and maintained at the Laboratory of Department of Pharmacy of Duy Tan University, Da Nang, Vietnam. For the assay (20,21), aliquots of *Syzygium* essential oils, dissolved by DMSO (1% stock solution), were placed in a 300-mL beaker and added to distilled water, containing 20 larvae (3th and early 4th instars). In each experiment, a set of controls using DMSO was also run for comparison. Mortality was calculated after 24 h and 48 h of exposure during which no nutritional supplement was added. The experiments have been carried out at room temperature. Each test was carried out with 3 replicates with several concentrations (100, 50, 25, 12.5, 6.0, 3.0, 1.5, and 0.75 µg/mL). Permethrin with the same tested concentrations was used as a positive control. The acute larvicidal effects on *Ae. aegypti* were recorded for 24 h and 48 h treatments. The data obtained were subjected to log-probit analysis to obtain LC₅₀ values, LC₉₀ values, and 95% confidence limits using XLSTAT v. 2018.5 (Addinsoft, Paris, France).

3. Results and discussion

A total of 46 compounds were identified in the leaf essential oil of *S. formosum*, which accounted for 94.1% (Table 1). This leaf essential oil was dominated by monoterpene hydrocarbons (51.1%), sesquiterpene hydrocarbons (27.0%), and oxygenated sesquiterpenes (15.2%). Oxygenated monoterpenes presented in a trace amount of 0.8%. The principal compounds were monoterpene hydrocarbons (*E*)- β -ocimene (18.0%) and β -pinene (16.8%), and sesquiterpene hydrocarbon bicyclogermacrene (9.1%). There were lesser percentages of *cis*- β -elemene (7.6%), spathulenol (6.6%), α -pinene (6.3%), (*Z*)- β -ocimene (3.8%), β -caryophyllene (3.2%), myrcene (2.4%), viridiflorol (2.1%), alismol (1.8%), germacrene D (1.6%), limonene (1.4%), and cubeban-11-ol (1.1%). The remaining compounds were observed with amounts of less than 1.0%.

Similar to *S. formosum* leaf essential oil, the extraction of *S. syzygioides* leaves gave a yellow essential oil with 47 identified compounds (94.9%), which was predominated by monoterpene hydrocarbons (39.6%) and sesquiterpene hydrocarbons (37.8%). Oxygenated sesquiterpenes were also significant with 16.6%, whereas oxygenated monoterpenes was present in a trace amount of 0.9%. α -Pinene (15.2%), (*Z*)- β -ocimene (10.3%), and

(*E*)- β -ocimene (7.1%) can be seen as characteristic monoterpene hydrocarbons. The principal sesquiterpene hydrocarbon, once again, was bicyclogermacrene (16.5%). Spathulenol (7.4%) was the most abundant compound in the group of oxygenated sesquiterpenes. Obviously, the amounts of α -pinene, bicyclogermacrene, and (*Z*)- β -ocimene in *S. syzygioides* leaf essential oil were higher than those in *S. formosum* leaf essential oil by 8.9, 7.4, and 6.5%, respectively. However, in contrast to *S. formosum* leaf essential oil, β -pinene (0.9%) and *cis*- β -elemene (1.1%) have been identified as minor compounds in *S. syzygioides* leaf essential oil. Likewise, the highest amount compound, (*E*)- β -ocimene in *S. formosum* leaf oil was associated with only 7.1% in *S. syzygioides* leaf essential oil.

The yellow essential oil of *S. megacarpum* leaf was obtained with 31 identified compounds (94.2%) that was almost entirely sesquiterpene hydrocarbons (83.8%) in character. In the meantime, their oxygenated derivatives reached 9.9%. Monoterpene hydrocarbons were not prevalent with 0.3%, while phenyl ethyl hexanoate (0.2%) represented as a non-terpenic compound only. The percentage of major compound bicyclogermacrene (37.0%) was recorded to outnumber the ones in the leaf essential oils of *S. formosum* and *S. syzygioides* by 27.9 and 20.5%, respectively. The percentages of β -caryophyllene and germacrene D in the leaf essential oils of *S. formosum* and *S. syzygioides* are not remarkable, but they achieved significant amounts of 9.4 and 16.4% in the leaf essential oils of *S. megacarpum*, respectively. Likewise, with 11.2%, *cis*- β -elemene in *S. megacarpum* leaf essential oil was found to outstrip that in the two first essential oils. *S. megacarpum* leaf oil also contained lower levels of spathulenol (3.6%), caryophyllene oxide (2.3%), δ -elemene (2.0%), α -humulene (1.9%), and δ -cadinene (1.0%).

Considering *S. chantaranothaianum* species, 36 compounds were identified in its leaf essential oil, which represented 96.2%. In the same manner with *S. megacarpum* leaf essential oil, sesquiterpene hydrocarbons (83.0%) were predominant in this oil sample, followed by oxygenated sesquiterpenes (12.2%), and monoterpene hydrocarbons (1.0%). Resembling *S. megacarpum* leaf essential oil, oxygenated monoterpenes were completely absent. The leaf essential oil of *S. chantaranothaianum* was associated with the appearance of two major components β -caryophyllene (19.4%) and bicyclogermacrene (24.7%). In addition, there have been accompanied by smaller percentages, such as germacrene D (7.9%), δ -elemene (7.5%), *cis*- β -elemene (5.4%), germacrene B (4.3%), and γ -elemene (3.1%), δ -selinene (1.6%), 9-*epi*-(*E*)-caryophyllene (1.3%), and aromadendrene and α -humulene (1.1%).

Table 1. The Identified Compounds (%) in the Essential Oils of Four *Syzygium* Leaves.

No.	R _E	R _L	Constituents	<i>S. formosum</i>	<i>S. syzygioides</i>	<i>S. megacarpum</i>	<i>S. chantaranthaianum</i>
1.	930	924	α -Thujene	0.3	0.3		
2.	939	932	α -Pinene	6.3	15.2		
3.	978	969	Sabinene	0.8	0.1		
4.	984	974	β -Pinene	16.8	0.9	0.3	
5.	991	988	Myrcene	2.4	0.9		
6.	1010	1002	α -Phellandrene	0.1	2.1		
7.	1029	1022	<i>O</i> -Cymene	0.5	1.0		
8.	1033	1024	Limonene	1.4	0.7		
9.	1035	1025	β -Phellandrene	0.2			
10.	1038	1032	(<i>Z</i>)- β -Ocimene	3.8	10.3		
11.	1049	1044	(<i>E</i>)- β -Ocimene	18.0	7.1		1.0
12.	1063	1054	γ -Terpinene	0.1			
13.	1093	1086	Terpinolene	0.4	0.4		
14.	1101	1095	Linalool	0.3	0.4	0.2	
15.	1131	1128	<i>allo</i> -Ocimene		0.6		
16.	1185	1174	Terpinen-4-ol		0.1		
17.	1257	1253	Linalyl acetate	0.3	0.4		
18.	1290	1282	<i>trans</i> -Linalool oxide acetate	0.2			
19.	1347	1335	δ -Elemene	0.4	0.5	2.0	7.5
20.	1384	1372	Geranyl acetate	0.3			
21.	1388	1374	α -Copaene	0.2	0.2	0.5	
22.	1402	1389	<i>cis</i> - β -Elemene	7.6	1.1	11.2	5.4
23.	1424	1409	α -Gurjunene		0.3	0.2	0.3
24.	1436	1417	β -Caryophyllene	3.2	3.8	9.4	19.4
25.	1444	1434	γ -Elemene	0.2			3.1
26.	1445	1436	α - <i>trans</i> -Bergamotene		0.7		
27.	1449	1440	β -Gurjunene		0.1	0.4	
28.	1452	1442	α -Maalinene		0.2		
29.	1455	1443	Aromadendrene	0.3	3.6	0.7	1.1
30.	1459	1444	Selina-5,11-diene		0.4		0.2
31.	1460	1445	(<i>Z</i>)- β -Farnesene				
32.	1462	1449	Guaia-1 (5), 6-diene				0.3
33.	1470	1467	α -Humulene	0.9	1.1	1.9	1.1
34.	1475	1469	Striatene				
35.	1478	1470	9- <i>epi</i> -(<i>E</i>)-Caryophyllene	0.6	1.6	0.7	1.3
36.	1489	1478	γ -Muurolene		0.7	0.6	0.7
37.	1491	1480	Valecene				0.5
38.	1493	1482	α -Amorphene	0.2	0.3	0.2	0.4
39.	1497	1484	Germacrene D	1.6	1.1	16.4	7.9
40.	1503	1489	β -Selinene	0.6	1.6	0.8	0.3
41.	1505	1498	δ -Selinene				1.6
42.	1511	1499	Viridiflorene	0.4	1.0		0.1
43.	1513	1500	Bicyclodermacrene	9.1	16.5	37.0	24.7
44.	1521	1511	δ -Amorphene			0.2	0.6
45.	1528	1513	γ -Cadinene	0.2	0.4	0.3	0.3
46.	1535	1522	δ -Cadinene	0.9	0.9	1.0	1.4
47.	1552	1537	α -Cadiene	0.2			
48.	1559	1545	Selina-3,7 (11)-diene				0.5
49.	1568	1561	(<i>E</i>)-Nerodiol			0.6	
50.	1576	1566	Germacrene B	0.4	0.5	0.3	4.3
51.	1583	1570	Dendrolasin		1.2		
52.	1586	1575	Globulol			0.5	
53.	1587	1576	Palustrol	0.4	0.5		0.7
54.	1596	1580	Spathulenol	6.6	7.4	3.6	1.1
55.	1603	1582	Caryophyllene oxide			2.3	
56.	1603	1589	Viridiflorol	2.1	2.9		2.6
57.	1610	1591	Guaiol			0.2	
58.	1612	1595	Cubeban-11-ol	1.1	1.7		2.9
59.	1620	1600	Rosifoliol	0.4	0.5		0.5
60.	1620	1602	Ledol	0.4		0.3	0.4
61.	1629	1608	Humulene epoxide II			0.3	
62.	1645	1627	1- <i>epi</i> -Cubenol	0.3	0.2		1.9
63.	1649	1630	γ -Eudesmol				0.2
64.	1650	1640	Phenyl ethyl hexanoate			0.2	
65.	1655	1644	Alismol	1.8			
66.	1656	1638	<i>epi</i> - α -Cadinol	0.4	2.1	0.5	0.4
67.	1658	1640	<i>epi</i> - α -Muurolol	0.2			0.2
68.	1661	1644	α -Muurolol		0.2		
69.	1671	1652	α -Cadinol	0.8	0.6	0.9	0.7
70.	1674	1658	<i>neo</i> -Intermedeol	0.7	0.5	0.5	0.6

(Continued)

Table 1. (Continued).

No.	RI _E	RI _L	Constituents	<i>S. formosum</i>	<i>S. syzygioides</i>	<i>S. megacarpum</i>	<i>S. chantaranthaianum</i>
			Total	94.1	94.9	94.2	96.2
			Monoterpene hydrocarbons	51.1	39.6	0.3	1.0
			Oxygenated monoterpenes	0.8	0.9		
			Sesquiterpene hydrocarbons	27.0	37.8	83.8	83.0
			Oxygenated sesquiterpenes	15.2	16.6	9.9	12.2
			Non-terpene compounds			0.2	

RI_E: Retention indices relative to *n*-alkanes (C₇-C₃₀) on HP-5 MS column.

RI_L: Retention indices from NIST standard database.

Bold: Compounds with greater than 5%.

To date, there were several GC-MS analytical studies on Vietnamese *Syzygium* species. (*Z*)- β -Ocimene (20.3%) and caryophyllene oxide (13.2%) were characteristic compounds of *S. nervosum* leaf oil from Southern Vietnam (22). Another example is that β -caryophyllene (23.40%), bicyclogermacrene (21.23%), and (*Z*)- β -ocimene (10.61%) established as the main compounds in the leaf essential oil of *S. tsoongii*, also collected from North Central Coast region of Vietnam (23). Thereby, it might be possibly concluded that monoterpene hydrocarbons and sesquiterpene hydrocarbons seem to be characters of the essential oils of Vietnamese *Syzygium* plants.

The obtained oils have been subjected to antimicrobial activity against three Gram-positive bacteria *E. faecalis*, *S. aureus*, and *B. cereus*, three Gram-negative bacteria *E. coli*, *P. aeruginosa*, and *S. enterica*, and one yeast *C. albicans*. As shown in Table 2, four samples showed powerful antimicrobial activity against Gram-positive bacteria *S. aureus* and *B. cereus* since these leaf oils possessed the MIC values better than those of the positive control streptomycin. In another case, the leaf essential oils of *S. megacarpum* (MIC = 16 μ g/mL) and *S. syzygioides* (MIC = 32 μ g/mL) were comparable with streptomycin (MIC = 32 μ g/mL) against Gram-positive bacterium *E. faecalis*, while two remaining essential oils exerted the same MIC value of 64 μ g/mL.

Regarding the Gram-negative bacteria, the MIC values assigning to four essential oils against *P. aeruginosa* orderly run as *S. megacarpum* and *S. chantaranthaianum* (MIC = 32 μ g/mL) >

S. syzygioides (MIC = 64 μ g/mL) > *S. formosum* and streptomycin (MIC = 128 μ g/mL). However, all samples failed to inhibit *E. coli* and *S. enterica*. Of the yeast *C. albicans*, the leaf oils of *S. formosum* and *S. syzygioides* were associated with the MIC value of 16 μ g/mL, as compared with that of the standard cycloheximide (MIC = 32 μ g/mL). The leaf essential oil of *S. chantaranthaianum* induced an equivalent MIC value of 32 μ g/mL to that of cycloheximide, whereas the leaf essential oil of *S. megacarpum* was strongly active against *C. albicans* with the same MIC value of 64 μ g/mL. As can be seen, the leaf essential oils of *S. formosum* and *S. syzygioides* are better than two remaining essential oils in antimicrobial assay against *C. albicans*. It may be due to the abundance of monoterpene hydrocarbons in these two samples.

This is the first time that Vietnamese *Syzygium* plants have been subjected to antimicrobial assays. The obtained results match well with previous reports. For instance, the Gram-positive bacterium *Bacillus subtilis* was susceptible to the leaf essential oils of *S. aromaticum* and *S. polyanthum* (MIC = 31.25 μ g/mL), but these oils were inactive to *E. coli* (23). Essential oil of clove *S. aromaticum* showed remarkable antimicrobial activity against 20 different isolates of *Listeria monocytogenes* at both concentrations of 1 and 10 μ g/mL growth medium, and inhibited *Streptococcus mutans* with the MIC value of 30 μ g/mL (24,25).

The 24-h and 48-h larvicidal activity of four *Syzygium* leaf essential oils against mosquito *Aedes aegypti* is shown in Table 3. It noted that essential oils have been considered actively larvicidal action if the

Table 2. Antimicrobial Activity of *Syzygium* Leaf Oils.

Samples	Gram (+)			Gram (-)			Yeast
	<i>E. faecalis</i>	<i>S. aureus</i>	<i>B. cereus</i>	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>S. enterica</i>	<i>C. albicans</i>
<i>S. formosum</i> leaf oil	64	64	32	-	128	-	16
<i>S. syzygioides</i> leaf oil	32	64	32	-	64	-	16
<i>S. megacarpum</i> leaf oil	16	64	32	-	32	-	64
<i>S. chantaranthaianum</i> leaf oil	64	32	32	-	32	-	32
Streptomycin	32	128	256	256	128	256	-
Cycloheximide	-	-	-	-	-	-	32

"-": inactive.

Table 3. Mosquito Larvicidal Activity of *Syzygium* Leaf Oils Against *Ae. Aegypti*.

Samples	LC ₅₀ (95% confidence levels)	LC ₉₀ (95% confidence levels)	χ^2	<i>p</i>
24-h treatment				
<i>S. formosum</i> leaf oil	30.1829 (28.1738–33.4702)	38.8101 (34.6692–47.9616)	0.0405804	0.980
<i>S. syzygioides</i> leaf oil	40.0409 (35.9258–43.1261)	50.6895 (47.2371–55.5091)	0.0178952	0.999
<i>S. megacarpum</i> leaf oil	28.3124 (26.4661–30.5315)	39.0577 (35.3101–46.0244)	0.0512473	0.975
<i>S. chantaranthaianum</i> leaf oil	27.5695 (25.4163–30.0059)	43.0427 (38.3092–51.1411)	6.11088	0.106
Permethrin (control)	0.0094 (0.0082–0.0107)	0.0211 (0.0185–0.0249)	57.6	0.000
48-h treatment				
<i>S. formosum</i> leaf oil	25.5460 (24.0321–27.6987)	33.0007 (30.1325–38.8792)	0.0010657	0.999
<i>S. syzygioides</i> leaf oil	39.2594 (35.1459–42.4376)	49.2314 (45.7001–53.9207)	0.0105410	1.000
<i>S. megacarpum</i> leaf oil	26.7038 (24.9437–29.2504)	35.7358 (32.3128–42.5290)	0.440285	0.802
<i>S. chantaranthaianum</i> leaf oil	26.6563 (24.6262–28.9373)	40.9687 (36.5517–48.6822)	6.15491	0.104

LC₅₀ value is less than 100 µg/mL (20). Based on this criterion, all studied samples exhibited good larvicidal activity with the 24-h and 48-h LC₅₀ values of around 25–40 µg/mL, as well as the 24-h and 48-h LC₉₀ values of around 33–50 µg/mL. In detail, the leaf essential oils of *S. megacarpum* and *S. chantaranthaianum* having the respective 24-h LC₅₀ values of 27 and 28 µg/mL are better than those of *S. formosum* leaf oil (LC₅₀ = 30 µg/mL) and *S. syzygioides* leaf oil (LC₅₀ = 40 µg/mL). This can be explained by the high concentration of sesquiterpene hydrocarbons, especially in terms of bicyclogermacrene. However, *S. formosum* leaf essential oil with the highest percentages of β -pinene (16.8%), (*E*)- β -ocimene (18.0%), and *cis*- β -elemene (7.6%) induced the lowest 48-h LC₅₀ value of 25 µg/mL, and the lowest 24-h and 48-h LC₉₀ values of 33–38 µg/mL. Significantly, our current results are superior to previous analogs. The leaf essential oils of Indian *S. lanceolatum* and Brazilian *S. jambolana* generated the LC₅₀ values of 55.11 and 433 µg/mL, respectively (26,27). The leaf essential oil of another Brazilian *S. aromaticum* caused the LC₅₀ value of > 90 µg/mL against *A. aegypti* (28). Thereby, it is expected that Vietnamese *Syzygium* essential oils are now promising agents for antimicrobial and mosquito repellency drug developments.

4. Conclusion

The current results exhibited a great variation in chemical compositions of four *Syzygium* leaf essential oils. Monoterpene hydrocarbons (39.6–51.1%) and sesquiterpene hydrocarbons (27.0–37.8%) represented the leaf essential oils of *S. formosum* and *S. syzygioides*, whereas the leaf essential oils of *S. megacarpum* and *S. chantaranthaianum* were dominated by sesquiterpene hydrocarbons (83.0–83.8%). Bicyclogermacrene (9.1–37.0%) was detected in four studied essential oils. β -Pinene (16.8%) and (*E*)- β -ocimene (18.0%) were also characteristics of *S. formosum* leaf oil, *S. syzygioides* leaf essential oil also contained α -pinene (15.2%), (*Z*)- β -ocimene

(10.3%), spathulenol (7.4%), and (*E*)- β -ocimene (7.1%). *cis*- β -Elemene (5.4–11.2%), β -caryophyllene (9.4–19.4%), and germacrene D (7.9–16.4%) were found in the leaf essential oils of *S. megacarpum* and *S. chantaranthaianum*. All the tested oils with the MIC values of 16–128 µg/mL were comparable or better than the positive control streptomycin against Gram-positive bacteria *S. aureus* and *B. cereus*, and Gram-negative bacterium *P. aeruginosa*. In the same manner, the leaf essential oils of *S. formosum*, *S. syzygioides*, and *S. chantaranthaianum* with the MIC values of 16–128 µg/mL were comparable or better than the positive control cycloheximide against the yeast *C. albicans*. Four oil samples have so far shown good larvicidal activity against mosquito *Ae. aegypti* with the 24-h and 28-h LC₅₀ values of 25–40 µg/mL, and the 24-h and 28-h LC₉₀ values of 33–50 µg/mL.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This research was funded by Ministry of Education and Training, Vietnam, under grant number: B2022-TDV-07.

ORCID

Le Duc Giang  <http://orcid.org/0000-0002-3269-9915>

References

1. R. Ascrizzi, L. Ceccarini, S. Tavarini, G. Flamini and L. G. Angelini, *Valorisation of hemp inflorescence after seed harvest: cultivation site and harvest time influence agronomic characteristics and essential oil yield and composition*. *Industrial Crops and Products*, **139**, 111541 (2019). doi: [10.1016/j.indcrop.2019.111541](https://doi.org/10.1016/j.indcrop.2019.111541).
2. P.S. Kakaraparthi, K.V.N.S. Srinivas, J.K. Kumar, A. N. Kumar, D.K. Rajput and V.U.M. Sarma, *Variation*

- in the essential oil content and composition of citronella (*Cymbopogon winterianus* Jowitt.) in relation to time of harvest and weather conditions. *Industrial Crops and Products*, **61**, 240–248 (2014). doi: [10.1016/j.indcrop.2014.06.044](https://doi.org/10.1016/j.indcrop.2014.06.044).
3. D.B. Marcum and B.R. Hanson, *Effect of irrigation and harvest timing on peppermint oil yield in California*. *Agricultural Water Management*, **82**(1–2), 118–128 (2006). doi: [10.1016/j.agwat.2005.07.022](https://doi.org/10.1016/j.agwat.2005.07.022).
 4. A. Carrubba and C. Catalano, *Essential Oil Crops for Sustainable Agriculture – A Review*. In Lichtfouse, E. (eds) *Climate Change, Intercropping, Pest Control and Beneficial Microorganisms*. Sustainable Agriculture Reviews, Vol. 2, Dordrecht. [10.1007/978-90-481-2716-0_8](https://doi.org/10.1007/978-90-481-2716-0_8)
 5. A.H. Hussein, A. Said-Al, A.S. Sabra, A. Alataway, T. Astatkie, A.A. Mahmoud and E. Bloem, *Biomass production and essential oil composition of thymus vulgaris in response to water stress and harvest time*. *Journal of Essential Oil Research*, **31**(1), 63–68 (2019). doi: [10.1080/10412905.2018.1518794](https://doi.org/10.1080/10412905.2018.1518794).
 6. W.K. Soh and J. Parnell, *A Revision of Syzygium Gaertn. (Myrtaceae) in Indochina (Cambodia, Laos and Vietnam)*. *Adansonia*, **37**(2), 179–275 (2015). doi: [10.5252/a2015n2a1](https://doi.org/10.5252/a2015n2a1).
 7. D.F. Cortés-Rojas, C.R.F. Souza and W.P. Oliveira, *Clove (Syzygium aromaticum): A precious spice*. *Asian Pacific Journal of Tropical Biomedicine*, **4**(2), 90–96 (2014). doi: [10.1016/S2221-1691\(14\)60215-X](https://doi.org/10.1016/S2221-1691(14)60215-X).
 8. S. Srivastava and D. Chandra, *Pharmacological potentials of Syzygium cumini: a review*. *Journal of the Science of Food and Agriculture*, **93**(9), 2084–2093 (2013). doi: [10.1002/jsfa.6111](https://doi.org/10.1002/jsfa.6111).
 9. P.M. Shafi, M.K. Rosamma, J. Jamil and P.S. Reddy, *Antibacterial activity of Syzygium cumini and Syzygium travancoricum leaf essential oils*. *Fitoterapia*, **73**(5), 414–416 (2002). doi: [10.1016/S0367-326X\(02\)00131-4](https://doi.org/10.1016/S0367-326X(02)00131-4).
 10. T.H. Khanh, P.H. Ban and T.M. Hoi, *Chemical composition of essential oils from the leaves of Syzygium bullockii and Syzygium tsoongii in ke go nature reserve, Ha Tinh province*. *VNU Journal of Science*, **37**(2), 18–23 (2020). doi: [10.25073/2588-1140/vnunst.4999](https://doi.org/10.25073/2588-1140/vnunst.4999).
 11. R. Sarvesan, P. Eganathan, J. Saranya and P. Sujanalal, *Chemical composition and antimicrobial activity of leaf essential oil of Syzygium grande (Wight) Walp.* *Journal of Essential Oil-Bear Plants*, **18**(3), 642–646 (2015). doi: [10.1080/0972060X.2014.958572](https://doi.org/10.1080/0972060X.2014.958572).
 12. A.K. Alanazi, M.H. Alqasbi, M. Alrouji, F.A. Kuriri, Y. Almuhanna, B. Joseph and M. Asad, *Antibacterial activity of Syzygium aromaticum (clove) bud oil and its interaction with imipenem in controlling wound infections in rats caused by methicillin-resistant staphylococcus aureus*. *Molecules*, **27**(23), 8551 (2022). doi: [10.3390/molecules27238551](https://doi.org/10.3390/molecules27238551).
 13. M. Govindarajan and G. Benelli, *α -humulene and β -elemene from Syzygium zeylanicum (myrtaceae) essential oil: highly effective and eco-friendly larvicides against Anopheles subpictus, Aedes albopictus, and Culex tritaeniorhynchus (Diptera: Culicidae)*. *Parasitology Research*, **115**(7), 2771–2778 (2016). doi: [10.1007/s00436-016-5025-2](https://doi.org/10.1007/s00436-016-5025-2).
 14. M. Osanloo, M.M. Sedaghat, F. Esmaeili and A. Amani, *Larvicidal activity of essential oil of Syzygium aromaticum (clove) in comparison with its major constituent, eugenol, against anopheles stephensi*. *Journal of Arthropod-Borne Diseases*, **12**, 361–369 (2018). doi: [10.18502/jad.v12i4.354](https://doi.org/10.18502/jad.v12i4.354).
 15. S.N. The, T.A. Le, D.T.T. Thuy, N.D. Luyen and T. T. Tuyen, *Essential oils from the aerial part and rhizome of amomum muricarpum Elmer and their antimicrobial activity*. *Letters in Applied NanoBioscience*, **11**, 3322–3328 (2021). doi: [10.33263/LIANBS111.33223328](https://doi.org/10.33263/LIANBS111.33223328).
 16. S.N. The, L.A. Tuan, T.D.T. Thu, N.D. Luyen and T. T. Thi, *Essential oils of the Asteraceae plants blumea riparia DC. And pluchea pteropoda Hemsl. ex Hemsl. growing in Vietnam*. *Natural Product Communications*, **17**(6), 1–6 (2022). doi: [10.1177/1934578X221110662](https://doi.org/10.1177/1934578X221110662).
 17. L.T. Huong, D.N. Dai, L.T.M. Chau and I. A. Ogunwande, *Analysis of Essential Oils from Alpinia napoensis*. *Chemistry of Natural Compounds*, **2018**, **54** (5), 992–994. doi: [10.1007/s10600-018-2532-7](https://doi.org/10.1007/s10600-018-2532-7).
 18. L.T. Huong, D.T.M. Chau, N.T.G. An, D.N. Dai and I. A. Ogunwande, *Essential oils of Lauraceae: antimicrobial activity and constituents of phoebe macrocarpa CY wu leaf essential oil from Vietnam*. *Journal of Essential Oil Bearing Plants*, **25**(2), 297–304 (2022). doi: [10.1080/0972060X.2022.2072694](https://doi.org/10.1080/0972060X.2022.2072694).
 19. L.T. Huong, L.D. Linh, D.N. Dai and I.A. Ogunwande, *Chemical compositions and antimicrobial activity of essential oils from Amomum velutinum X.E.Ye, Škornièk. & N.H.Xia (Zingiberaceae) from Vietnam*. *Journal of Essential Oil-Bear Plants*, **23**, 1132–1141 (2020). doi: [10.1080/0972060X.2020.1856005](https://doi.org/10.1080/0972060X.2020.1856005).
 20. N.H. Hung, N.T. Huong, N.T. Chung, N.T.H. Thuong, P. Satyal N.A. Dung, et al., *Callicarpa species from central Vietnam: Essential oil compositions and mosquito larvicidal activities*. *Plants*, **2022**, **9**(1), 113. doi: [10.3390/plants9010113](https://doi.org/10.3390/plants9010113).
 21. E.F.L. Cardoso Bailao, D. Godoi Pereira, C. Aline Romano, A.T.D. Santana Paz, T.M.E. Silva J.R. D. Paula, et al., *Larvicidal effect of the citrus limettoides peel essential oil on Aedes aegypti*. *South Africa Journal of Bot*, **2022**, **144**, 257–260. doi: [10.1016/j.sajb.2021.09.013](https://doi.org/10.1016/j.sajb.2021.09.013).
 22. N.T.G. An, L.T. Huong, P. Satyal, T.A. Tai, D.N. Dai N. H. Hung, et al., *Mosquito larvicidal activity, antimicrobial activity and chemical compositions of essential oils from four species of Myrtaceae from central Vietnam*. *Plants*, **2020**, **9**(4), 544. doi: [10.3390/plants9040544](https://doi.org/10.3390/plants9040544).
 23. A. Hamad, M.G.P. Mahardika, I. Yuliani and D. Hartanti, *Chemical constituents and antimicrobial activities of essential oils of Syzygium polyanthum and Syzygium aromaticum*. *Rasayan Journal of Chemistry*, **10**, 564–569 (2017). doi: [10.7324/RJC.2017.1021693](https://doi.org/10.7324/RJC.2017.1021693).
 24. S.G. Deans, R.C. Noble, R. Hiltunen, W. Wuryani and L.G. Penzes, *Antimicrobial and antioxidant properties of Syzygium aromaticum (L.) merr. & Perry: impact upon bacteria, fungi and fatty acid levels in ageing mice*. *Journal of Flavour Fragrance*, **10**, 323–328 (1995). doi: [10.1002/ffj.2730100507](https://doi.org/10.1002/ffj.2730100507).
 25. K. Koba, A.Y. Nenonene, C. Raynaud, J.P. Chaumont and K. Sanda, *Antibacterial activities of the buds essential oil of Syzygium aromaticum (L.) merr. & Perry from*

- Togo*. Journal of Biological Active Products and Nature, **1**(1), 42–51 (2011). doi: [10.1080/22311866.2011.10719072](https://doi.org/10.1080/22311866.2011.10719072).
26. E.S.B. Cavalcanti, S.M.D. Morais, M.A.A. Lima and E. W.P. Santana, *Larvicidal activity of essential oils from Brazilian plants against Aedes aegypti L.* Memórias do Instituto Oswaldo Cruz, **99**(5), 541–544 (2004). doi: [10.1590/s0074-02762004000500015](https://doi.org/10.1590/s0074-02762004000500015).
27. G. Benelli, M. Rajeswary and M. Govindarajan, *Towards green oviposition deterrents? Effectiveness of Syzygium lanceolatum (Myrtaceae) essential oil against six mosquito vectors and impact on four aquatic biological control agents.* Environmental Science and Pollution Research, **25**(11), 10218–10227 (2016). doi: [10.1007/s11356-016-8146-3](https://doi.org/10.1007/s11356-016-8146-3).
28. A.F.D. Oliveira Araujo, J.T. Ribeiro-Paes, J.T.D. Deus, S.C.D. Holanda Cavalcanti, R.D. Souza Nunes P. B. Alves, et al., *Larvicidal activity of Syzygium aromaticum (L.) merr and Citrus sinensis (L.) osbeck essential oils and their antagonistic effects with temephos in resistant populations of Aedes aegypti.* Memórias do Instituto Oswaldo Cruz, **2016**, **111**(7), 443–449. doi: [10.1590/0074-02760160075](https://doi.org/10.1590/0074-02760160075).