



**HỘI KHOA HỌC KỸ THUẬT PHÂN TÍCH HÓA, LÝ VÀ SINH HỌC VIỆT NAM**  
**VIETNAM ANALYTICAL SCIENCES SOCIETY**

ISSN - 0868 - 3224

**Tạp chí**  
**PHÂN TÍCH**  
**HÓA , LÝ VÀ SINH HỌC**  
*Journal of Analytical Sciences*

**T - 25**

**Số 2**

---

**2020**

**HA NOI**

- 31 Ứng dụng chiết điểm mù sử dụng 1-(2-pyridylazo)-2-naphthol và triton x-100 phân tích dạng mangan trong mẫu chè 177  
Application of cloud point extraction use 1- (2-pyridylazo) -2-naphthol and triton x-100 speciation analysis of manganese in tea samples  
*Lê Sỹ Bình, Đào Văn Bảy, Vũ Đức Lợi*
- 32 Polybrom diphenyl ete trong mẫu bụi nhà tại khu vực Hà Nội: phân tích và đánh giá 185  
Polybrominated diphenyl ethers in house dust from urban areas in Hanoi: analysis and assessment  
*Lê Trường Giang, Phạm Quốc Trung, Trịnh Thu Hà*
- 33 Tầm soát các hóa chất bảo vệ thực vật trong đất ruộng lúa bị ngập lụt ở miền Trung Việt Nam 191  
Screening pesticides in flooded paddy soil in center Vietnam  
*Lê Trường Giang, Dương Thị Hạnh, Trịnh Thu Hà*
- 34 Đánh giá khả năng quang xúc tác phân hủy p,p' DDT sử dụng TiO<sub>2</sub> phủ trên hạt silica gel 198  
Determination the photocatalytic activity in degradation P, P' ddt using TIO<sub>2</sub>immobilized on silica gel beads  
*Nguyễn Thị Huệ, Hà Thanh Hòa, Nguyễn Mạnh Nghĩa*
- 35 Ứng dụng Apatit / TiO<sub>2</sub> pha tạp nitơ để khử màu methylene xanh (MB) và xử lý Benzene, Toluene và Xylene (BTX) trong không khí 203  
Application of apatite coating on TiO<sub>2</sub> doped N for decoloring methylene blue (MB) and treating benzene, toluene and xylene (BTX) in the air  
*Nguyễn Thị Huệ*
- 36 Xác định đồng thời Aspartame và Saccharin trong một số loại đồ uống bằng phương pháp sắc ký lỏng hiệu năng cao 207  
Simultaneous identification of aspartame and saccharin in some daily beverages by using high performance liquid chromatography method  
*Dương Thị Tú Anh, Phạm Thanh Tùng, Đoàn Mạnh Cường*
- 37 Optimization of total phenolic and total flavonoid extraction conditions from leaves of launaea sarmentosa using the response surface methodology 214  
*Nguyen Tan Thanh, Tran Dinh Thang, Nguyen Thi Huyen, Tran Thi Thu Ha*
- 38 Synthesis and properties of complexes of some heavy rare earth ions and benzoic acid 221  
*Nguyen Thi Hien Lan, Mai Xuân Truong, Tran Thi Phuong*
- 39 Synthesis of some extended heterocyclic pyrazolo-pyridazine derivatives bearing (methyl)phenyl amide moiety 226  
Tổng hợp một vài dị vòng pyrazolo-pyridazine có chứa phần (methyl)phenyl amide  
*Tran Quang De, Nguyen Trong Tuan, Bui Thi Bui Hue*
- 40 Synthesis of some potential semi-conductor compounds via arylation of benzo[1,2-b:4,5-b']dithiophene 232  
*Tran Thi Phuong, Nguyen Thi Van Khanh, Nguyen Hien, Duong Quoc Hoan*
- 41 NMR and X-ray crystal structure analysis of arylated thieno[3,2-b]thiophene by C<sub>sp</sub><sup>2</sup>-H functionalization 237  
*Nguyen Hien, Nguyen Bich Ngan, Duong Quoc Hoan*
- 42 Photocatalytic degradation of methylene blue dye by zinc oxide nanoparticles obtained from green method 245  
Nghiên cứu hoạt tính quang xúc tác phân hủy methylen xanh của vật liệu ZnO kích thước nano tổng hợp theo phương pháp hóa học xanh  
*Nguyen Ngoc Thinh, Nguyen Van Anh, Nguyen Thi Anh Huong*

## OPTIMIZATION OF TOTAL PHENOLIC AND TOTAL FLAVONOID EXTRACTION CONDITIONS FROM LEAVES OF LAUNAEA SARMENTOSA USING THE RESPONSE SURFACE METHODOLOGY

Đến tòa soạn 28-11-2019

**Nguyen Tan Thanh, Tran Dinh Thang, Nguyen Thi Huyen**

*School of Chemistry, Biology and Environment, Vinh University*

**Tran Thi Thu Ha**

*School of natural Sciences Education, Vinh University*

*Tourism and Hospitality Department, Ha Tinh College of Technology*

The Central composite design (CCD) of response surface methodology (RSM) was used to investigate the effects of three factor as extraction temperature ( $^{\circ}\text{C}$ ), extraction time (min) and ethanol concentration (%) of *Launaea sarmentosa* leaves on the responses: total phenolic content (TPC) and total flavonoid content (TFC). The optimal conditions obtained from response RSM were 90% v/v for the solvent ratio,  $54^{\circ}\text{C}$  for extraction temperature and 110min for extraction time. The experimental values of TPC and TFC were  $318.85 \pm 0.32$  mgGAE/g,  $8.21 \pm 0.14$  mgCE/g.

**Keywords:** *Launaea sarmentosa*, total phenolic content (TPC), total flavonoid content (TFC), response surface methodology (RSM), extraction.

### 1. INTRODUCTION

*Launaea sarmentosa* (Willd) Schultz-Bip.ex Kuntze, belongs to family of Asteraceae, in Vietnam, It is grown at sandy coasts of Thai Binh, Nghe An, Ha Tinh, Ben Tre, Quang Tri [1], it is a creeping herb, native to tropical Indian coastlines. All parts of the *Launaea sarmentosa* (Willd.) plant especially leaves of contain high amounts of phenolic and flavonoid compounds with potential antioxidant properties. This Plant also synthesize huge amount of aromatic compound among which phenols or their oxygen substituted derivatives are predominant. These compounds provide protection against microbes for the plant [2].

*Launaea* has great importance due to its ethnobotanics, phytochemistry and biological activity, and various secondary metabolites including sesquiterpenoids, terpenoids and flavonoids [3]. It's root contains the following chemical components: calcium oxalate crystals, tannin content, contains alkaloids,

aminoacids, carbohydrates, glycosides, tannin, and steroids [4].

The role of flavonoids is to be the "biochemical repairman of nature", helping to correct errors for metabolic reactions, the biosynthesis processes of living ingredients, supporting endocrine regulation. Flavonoids are class of secondary plant metabolites with significant antioxidant and chelating properties. Antioxidant activity of flavonoids depends on the structure and substitution pattern of hydroxyl groups [5].

Response surface methodology (RSM) is an effective statistical method for optimizing experimental conditions and investigation of critical processes as well as reducing the number of experimental trials. RSM helps to define effects of the independent variables, whether it is alone or combination in the process [6,7]. One of the most important points in the implementation of this method is that the predicted values in the model should be verified experimentally. Thus, RSM is a useful



tool for optimizing the technology process over the conventional one factor at a time approach, which is relatively expensive and time-consuming. In this study, we have optimized the extraction conditions of total phenolic and total flavonoid from leaves of *Launaea sarmentosa* because these are two compounds found very much in genus *launaea* [8].

## 2. MATERIAL AND METHODS

### 2.1. Material

Leaves of *Launaea sarmentosa* were collected in Nghi Xuan District of Ha Tinh Province, Vietnam in September 2019 and identified by Institute of Ecology and Biological Resources, Vietnam Academy of Science and Technology. A voucher specimen was deposited at the herbarium of the School of Chemistry, Biology and Environment, Vinh University. The material is dried, crushed and stored at 4°C for further experiments.

### 2.2. Methods

#### 2.2.1. Total Phenolic Content (TPC)

The TPC of the *Launaea sarmentosa* leaves extracts was measured according to the method reported by Singleton et al. [9] with a little modification. This method is based on measuring color change caused by reagent by phenolates in the presence of sodium carbonate. 1ml of sample was mixed with 5ml of Folin-Ciocalteu's solution. After 3 min, 4ml of 7.5% sodium carbonate solution was added to a mixture and adjusted to 10ml with deionized water. The mixture was kept at room temperature in a dark environment for 60min. The color change was determined by scanning the wavelength at 765nm (Agilent 8453 UV – Visible Spectrophotometer) since maximum absorbance was obtained. TPC of the *Launaea sarmentosa* leaves extract was determined as mg gallic acid equivalent using the standard curve prepared at different concentrations of gallic acid and reported as mgGAE/g dry weight (DW).

#### 2.2.2. Total Flavonoid Content (TFC)

The TFC of the *Launaea sarmentosa* leaves extract was estimated according to the procedures described by D. Marinova et al.[10]

with slight modification. An aliquot (1ml) of extracts or standard solution of catechin (0.01 ÷ 0.07mg/ml) was added to 10 volumetric flask containing 4 ml of dd H<sub>2</sub>O. To the flask was added 0.3ml 5%NaNO<sub>2</sub>. After 5 min, 0.3ml 10% AlCl<sub>3</sub> was added. At 6<sup>th</sup> min, 2ml 1M NaOH was added and the total volume was made up to 10ml with ddH<sub>2</sub>O. The solution was mixed well and the absorbance was measured against prepared reagent blank at 510nm (Agilent 8453 UV-Visible Spectrophotometer). Total flavonoid content of *Launaea sarmentosa* leaves extract was expressed as mg Catechin equivalents mgCE/g DW.

#### 2.2.3. Experimental design

Before the development of the study by RSM, determination of experimental ranges for independent variables namely extraction time, extraction temperature, solvent/material ratio and ethanol concentration were carried out using total phenolic content as a determinant factor. Then, RSM was used to determine the optimum levels of extraction time (min), temperature (°C) and ethanol concentration (%) as extraction medium on two responses TPC and TFC in the *Launaea sarmentosa* leaves extracts. These three factors, namely extraction temperature (X<sub>1</sub>), extraction time (X<sub>2</sub>) and ethanol concentration (X<sub>3</sub>) were coded into three levels (-1, 0, +1). Ranges of extraction temperature, extraction time and ethanol concentration and the central point were selected based on preliminary experimental results. Statistical analysis on the means of triplicate experiments was carried out using the ANOVA procedure of the design expert software, version 7.0.

## 3. RESULTS AND DISCUSSION

### 3.1. Fitting the response surface models

The responses consisting of total phenolic content and total flavonoid content for *Launaea sarmentosa* leaves extract were optimized based on the central composite design (CCD), the CCD was used to identify the relationship between the response functions and process variables as well as to find out the

conditions that optimized the extraction process. The experimental design and corresponding three response variables are presented in Table 1. This design consisted of 20 experimental points with six replicates at the central point. In the present study, according to the sequential model sum of

squares, the highest order polynomials were utilized to select the models where the additional coefficients estimates were significant and the models are not aliased. Hence, for all three independent variables and responses, a quadratic polynomial model was selected and fitted well as suggested by the software.

Table 1: The experimental data obtained for the three responses based on the CCD matrix

RUN	X <sub>1</sub> (°C)	X <sub>2</sub> (min)	X <sub>3</sub> (%)	TPC Y <sub>1</sub> (mgGAE/g)	TFC Y <sub>2</sub> (mgCE/g)
1	50	120	90	318.15	8.32
2	60	100	80	318.05	8.18
3	60	100	80	318.48	8.17
4	70	120	70	318.76	8.01
5	60	100	80	318.16	8.17
6	76.82	100	80	331.61	7.95
7	43.18	100	80	311.98	8.03
8	60	100	96.82	315.01	8.29
9	60	66.36	80	311.16	8.25
10	60	133.64	80	318.11	8.21
11	50	80	70	294.65	8.02
12	70	80	70	317.65	8.15
13	60	100	80	318.08	8.15
14	50	120	70	312.62	7.93
15	70	120	90	318.27	8.16
16	70	80	90	327.83	8.12
17	60	100	80	318.39	8.16
18	60	100	63.18	300.93	7.99
19	60	100	80	318.09	8.17
20	50	80	90	311.67	8.27

The values of the two evaluation indices for each extracting condition were listed in Table 1. At extracting condition: 76.82°C, 80% ethanol concentration in 100min, the maximal TPC was 331.61 mgGAE/g and the maximal TFC was 8.32 mgCE/g at 50°C, 90% ethanol concentration in 120 min.

The final empirical regression model of their relationship between responses and the three tested variables for phenolic and flavonoid contents could be expressed by the following quadratic polynomial equation [Eqs. (1-2)]:

$$Y_1 = 318.20 + 5.74X_1 + 2.03X_2 + 4.09X_3 - 4.11X_1X_2 - 1.61X_1X_3 - 2.27X_2X_3 + 1.34X_1^2 - 1.19X_2^2 - 3.54X_3^2 \quad (1)$$

$$Y_2 = 8.17 - 0.017X_1 - 0.015X_2 + 0.093X_3 - 0.065X_1X_3 + 0.04X_2X_3 - 0.061X_1^2 + 0.024X_2^2 - 0.008X_3^2 \quad (2)$$

Where Y<sub>1</sub> is total phenolic content, Y<sub>2</sub> is the total flavonoid content, X<sub>1</sub> is the temperature, X<sub>2</sub> is the time and X<sub>3</sub> is the solvent ratio (ethanol concentration ratio).



Table 2: Analysis of variance (ANOVA) for the model

Source	Y <sub>1</sub> – Total phenolic content			Y <sub>2</sub> – Total flavonoid content		
	Mean Square	F- value	p- value	Mean Square	F- value	p-value
Model	132.24	1586.23	< 0.0001***	0.027	181.35	< 0.0001***
X <sub>1</sub> (temperature)	450.45	5403.24	< 0.0001***	0.004	27.55	0.0004***
X <sub>2</sub> (time)	56.136	673.36	0.0001***	0.003	21.52	0.0009***
X <sub>3</sub> (solvent ratio)	228.96	2746.50	0.0001***	0.117	800.83	< 0.0001***
X <sub>1</sub> X <sub>2</sub>	135.30	1622.94	< 0.0001***	4.5E-004	3.08	0.1099 <sup>NS</sup>
X <sub>1</sub> X <sub>3</sub>	20.672	247.97	< 0.0001***	0.034	231.18	< 0.0001***
X <sub>2</sub> X <sub>3</sub>	61.38	736.29	0.0001***	0.013	87.55	< 0.0001***
X <sub>1</sub> <sup>2</sup>	26.00	311.95	0.0001***	0.054	370.82	< 0.0001***
X <sub>2</sub> <sup>2</sup>	20.34	244.00	0.0001***	0.008	54.51	< 0.0001***
X <sub>3</sub> <sup>2</sup>	181.06	2171.79	< 0.0001***	9.94E-004	6.80	0.0262*
Lack of Fit	0.13	4.06	0.0751 <sup>NS</sup>	1.857E-004	1.74	0.2787 <sup>NS</sup>
R <sup>2</sup>	0.9993			0.9939		
C.V%	0.091			0.15		

\*p< 0.05; \*\*p< 0.01; \*\*\*p< 0.001; NS: non-significant.

The RSM model coefficients were validated by analysis of variance (ANOVA) of the response variables for the quadratic polynomial model summarized in Table 2. The ANOVA analysis results for multiple regression and response surface quadratic model of Y<sub>1</sub> and Y<sub>2</sub> were evaluated using the corresponding p and R<sup>2</sup> values. F values of Y<sub>1</sub> and Y<sub>2</sub> were calculated to be 1586.23 and 181.35, both leading to a p value <0.05, suggesting both the models were statistically significant. The models' coefficient of determination (R<sup>2</sup>) were 0.9993 and 0.9939, indicating that more than 99.93% and 99.39% of the response variability were explained, and supporting a good accuracy and ability of the established model within the range limits used. The F-values of Lack of Fit for Y<sub>1</sub> and Y<sub>2</sub> were 4.06 and 1.74, respectively,

implying that the Lack of Fit was not significant relative to the pure error. This indicated that the accuracy of the polynomial model was adequate.

### 3.2. Response surface analysis

Three factors that temperature, time and ethanol concentration effects the extraction condition of the maximum total phenolics and total flavonoids content. This section discusses how these conditions work on natural antioxidants extraction. Three-dimensional model graphs were plotted as shown in the respective figures. The response surface plots of the model were done by varying two variables, within experimental range under investigation and holding the other variables at its central level.

#### 3.2.1. Response surface analysis of total phenolic content




Figure 1: The response surface plot of TPC

The response surface plots for total phenolic extraction of *Launaea sarmentosa* leaves extract are shown in Fig. 1 demonstrating the effect and interaction of independent variables on the yields of total phenolics. As shown in Fig. 1 and Table 2, all of three factors (extraction temperature, extraction time and ethanol concentration ratio) have showed negative quadratic effects ( $p < 0.0001$ ). In fig. 1a, The surface plot demonstrates the function of extraction temperature versus time effect on TPC at fixed ethanol concentration (80%). We can be observed that the yields of total phenolic content increased with the increase of extraction temperature from 50°C to 70°C and the maximum amount of phenolics can be achieved at the highest temperature of 65±70°C at the shortest extraction time at 120 min. Higher solubility and diffusion coefficient of polyphenols were observed with increased temperature, allowing more extraction rate [11]. However, an upper limit of temperature must be respected in order to prevent decomposition of thermo sensitive phenolics during extraction [12]. These results are similar to a study reported by of Rajha et al. [13] which showed the total phenolics from grape by products increased with the increment of temperature and reduction of time.

The surface plot in Fig. 1b show the function of temperature versus solvent ratio effect on TPC at extraction time (120min). The yields of TPC increased with the increase of ethanol

concentration from 70%v/v to 90%v/v and the maximum phenolic content in *Launaea sarmentosa* leaves can be achieved at highest ethanol concentration (90%). The higher phenolic content could be explained by the natural polarity of the solvents used [14]. Ethanol and water were used in this study because they are safer to handle as compared to other organic solvents and more importantly, they are acceptable for human consumption. Samuagam et al. [15] stated that a suitable solvent ratio is able to improve the efficiency of extraction. The maximum total phenolic content in *Launaea sarmentosa* leaves can be obtained with optimum ethanol concentration and an extraction temperature of approximately 80±90 v/v% and 65±70°C respectively.

In Fig. 1c. The surface plots revealed that the higher TPC in *Launaea sarmentosa* leaves can be obtained when conducted at increasing ethanol concentration at fixed extraction time. Based on the result at constant extraction time of 120 min, 90% of ethanol concentrations yielded the most TPC as compared with 70% ethanol concentrations. These overall results of phenolic content indicate a similar trend as observed in the phenolic content in other study [16], [17], where the TP contents increased with increasing the independent variables ethanol concentration and processing time until a maximum amount of phenolic was reached

### 3.2.2. Response surface analysis of total flavonoid content




Figure 2: The response surface plot of TFC



The 3D in Fig.2a shows the response surface plot of temperature ( $X_1$ ) and time ( $X_2$ ) at fixed extraction solvent ratio (80%). Response surface plot showed that extraction temperature exhibited a weaker effect whereas extraction time represented a relatively significant effect on the favonoids yield. An increase in the yield of favonoid could be significantly achieved with the increase of extraction time, at any level of extraction temperature. Therefore, the optimum amount of favonoid was achieved in this study at 50±55°C and 110±120 min of extraction time.

The 3D surface plots in Fig. 2b shows the interaction between extraction temperature ( $X_1$ ) and solvent ratio ( $X_3$ ) at the fixed 100 min. According to Bazykina et al. [18] favonoids and their glycosides are thought to be efficiently extracted from plant materials by ethanol solvent. It was observed that the value of TFC in *Launaea sarmentosa* leaves increased when ethanol concentration was increased from 70 to 90v/v% at fixed 60°C extraction temperature. In contrast, increasing the extraction temperature at highest ethanol concentrations resulted to decreased, TFC values.

Fig. 2c shows the interaction between extraction time ( $X_2$ ) and ethanol concentration ( $X_3$ ) at the fixed extraction temperature at

60°C. An increase in ethanol concentration promoted the breakdown of the cell membrane that enhanced the permeability of the solvent into a solid matrix. In this study, highest favonoids content can be achieved when conducted at highest ethanol to water ratio 90% as compared with 30% with increasing extraction time. A great increase in the yield also resulted when extraction time was increased in the range of 80±120min.

### 3.3. Optimization and Model Verification

The final result for the simultaneous optimization using the desirability function approach suggested that the optimal ethanolic extraction conditions for *Launaea sarmentosa* leaves extract were at 54°C with 110 min and 90% of ethanol concentration to achieve the best combination for highest total phenolic and favonoids content. Table 3 shows the predicted and experimental values for the extraction of target compounds from *Launaea sarmentosa* leaves. The actual values obtained from the experimental gave the extraction yields of total phenolic and total flavonoid as 318.85±0.32mgGAE/g and 8.21±0.14mgCE/g. These experimental values were close to the predicted values (TPC = 320.53mgGAE/g, TFC = 8.23mgCE/g) derived from the respective regression models with the CV ranging from 0.24% to 0.52%.

Table 3: Comparison between the predicted and experimental values for antioxidants from extracts of *Launaea sarmentosa* leaves

Condition	Response values	
	Total phenolic content mgGAE/g	Total flavonoid content mgCE/g
Predicted	320.53	8.23
Experimental	318.85±0.32	8.21±0.14
% Difference (CV)	0.52%	0.24%

## 4. CONCLUSION

Use response surface methodology (RSM) with central composite design (CCD) were successfully developed to determine the optimum process parameters and the second order polynomial models for predicting responses were obtained. The best combination of extraction temperature, time and ethanol concentrations were found to be 54°C with 110 min and ethanol concentration ratio 90% which

rendered a mean phenolic content of 318.85±0.32 mgGAE/g and 6.12 ± 0.23 mgCE/g of total favonoid content from experimental run and thus indicated good antioxidant activities from the leaves of *Launaea sarmentosa*.

## REFERENCES

1. Thao H.X., Anh L.T, My N.T.D., *Đa dạng thực vật có hoa vùng đất cát huyện Triệu Phong và Hải Lăng tỉnh Quảng Trị*, The



- seventh national scientific conference on ecology and biological resources, 936-946 (2017).
2. Shalahuddin Millat. Md, Safiquil Islam, Saddam Hussain.Md, Mizanur Rahman Moghal.Md, Tarequl Islam., *Anti-bacterial Profiling of Launaea sarmentosa (willd.) and Brugiera cylindrical (L.): Two distinct ethno medicinal plants of Bangladesh.* Eur. J. Exp. Bio., **7**, 1-6 (2017).
  3. Abdelkrim C., Mebarka B., Nasser B. and Houria D., *Phytochemical and biological studies on Launaea Cas.genus (Asteraceae) from Algerian Sahara.* Cur. Top. Phyto., **11**, 67-80 (2012).
  4. Yusriyya S., Harischa C.R., Vinay J. S., Rabinarayan A., *Pharmacognostical evaluation of Launaea sarmentosa (willd.)schultz-bip.ex Kuntze root,* Ayu. J., **34**, 90-94 (2013).
  5. Milan S.Stankovie, *Total phenolic content, flavonoid concentration and antioxidant activity of Marrabiun peregrinum L.extracts.* Kragujevac J.Sci., **33**, 63-72 (2011).
  6. Bas D., Boyaci I.H., *Modeling and optimization I: usability of response surface methodology,* J. Food Eng., **788**, 36–845 (2007).
  7. Silva E.M., Rogez H., Larondelle Y., *Optimization of extraction of phenolics from Inga edulis leaves using response surface methodology,* Separ.Purif. Technol., **55**, 381–387 (2007).
  8. Le Thi Them, Phu Tuong Nguyen Dung, Pham Thi Nhat Trinh, Quach Tong Hung, Le Nguyen Tuong Vi, Nguyen Trong Tuan, Tri Duc Lam, Van Thuy Nguyen, Le Tien Dung, *Saponin, Polyphenol, Flavonoid content and  $\alpha$ -glucosidase Inhibitory Activity, Antioxidant Potential of Launaea sarmentosa Leaves grown in Ben Tre province, Vietnam,* Mat. Sci. Eng., **542**, 1-5 (2019).
  9. Singleton V.L., Orthofer R., Lamuela-Raventos R.M., *Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent,* Meth. Enzymol., **299**, 152-178 (1999).
  10. Marinova D., Ribarova F., Atanassova M., *Total phenolics and total flavonoid in Bulgarian fruits and vegetables,* J. Univ. Chem. Tech. Metallurg., **40(3)**, 255-260 (2005).
  11. Cacace J.E., Mazza G., *Mass transfer process during extraction of phenolic compounds from mixed berries,* J. Food Eng. **59**, 379–389 (2003).
  12. Silva E.M., Rogez H., Larondelle Y., *Optimization of extraction of phenolics from Inga edulis leaves using response surface methodology,* Separ. Purif. Technol., **55**, 381–387 (2007).
  13. Rajha H.N., Darra N.E., Hobaika Z., Boussetta N., Vorobiev E., Maraoun R.G., Louka N., *Extraction of total phenolic compound, favonoids, anthocyanins, and tannis from grape byproducts by response surface methodology. Influence of solid-liquid ratio, particle size, time, temperature and solvent mixtures on the optimization process.* Food Nutr. Sci., **5**, 397–409 (2014).
  14. Tan M.C., Tan C.P., Ho C.W., *Effects of extraction solvent system, time and temperature on total phenolic content of henna (Lawsonia inermis) stems.* Int. Food Res. J. **20**, 3117–3123 (2013).
  15. Samuagam L., Sia C.M., Akowuah G.A., Okechukwu P.N., Yim H.S., *The effect of extraction conditions on total phenolic content and free radical scavenging capacity of selected tropical fruits' peel.* Heal. Environ J. **4**, 80–102 (2013).
  16. Archana A., Bharathi V.D., Saraboji S., Thirunavukkarasu A., Nithya R., *Optimization and Extraction of Phenolic Compounds from Capsicum annum Using Response Surface Methodology ,* Int. J. Emerg. Res. Man. Tech. **4(7)**, 204-211 (2015).
  17. Aybastier O., Isik E., Sahin S., Demir C., (2013) *Optimization of ultrasonicassisted extraction of antioxidant compounds from blackberry leaves using response surface methodology.* Ind. Crop. Prods., **44**, 558–565.
  18. Bazykina N.I., Nikolaevskii A.N., Filippenko T.A., Kolerva V.G., *Optimization of conditions for the extraction of natural antioxidants from raw plant materials.* Pharm. Chem. J., **36**, 46–49 (2002).