

CONTENTS

		pages
1	Do Huu Nghi, Le Mai Huong, Le Huu Cuong, Do Huu Chi, Vu Dinh Giap, Nguyen Manh Cuong - Application of image-based high content analysis for the screening of bioactive natural-products.	1
2	Nguyen Thuy Chinh, Thach Thi Loc, Le Duc Giang, Ngo Phuong Thuy, Vu Thi Hien, Thai Hoang - Effect of polycaprolactone on characteristics and morphology of alginate/chitosan/lovastatin composite films.	13
3	Nguyen Dinh Tuan, Nguyen Phi Hung, Do Huu Nghi - Insulin-mimetic biflavones from a Vietnamese medicinal plant <i>Selaginella tamariscina</i> .	22
4	Dinh Thi Kim Hoa, Pham Quoc Long, Doan Lan Phuong - Research on the composition of lipids, fatty acids, and amino acids from egg and body of sea urchin <i>Tripneustes gratilla</i> .	30
5	Nguyen Thi Thuy Nga, Vu Thuy An, Dao Duy Quang - Antioxidant properties of folic acid: a DFT study.	39
6	Ngo Dang Truong Hai, Ngo Thi Chinh, Pham Minh Quan, Dao Duy Quang - Effects of substituents on C-H bond dissociation enthalpies of ent-kaurane diterpenoids: a DFT study.	46
7	Nguyen Chi Dung, Pham Thi My Ninh, Dinh Minh Hiep - A comparison of the cytotoxic activity of extracts from fruiting bodies and mycelial biomass of <i>Cordyceps neovolkiana</i> (DL004) fungus.	53
8	Do Thi Hoa Vien, Dao Thi Trang - Research on extraction of isoflavones from soybean germ.	61
9	Nguyen Phi Hung, Nguyen Dinh Tuan, Pham Quoc Long - Triterpenoids from <i>Selaginella rolandi-principis</i> and their PTP1B inhibitory activity.	69
10	Ha Bao Chau, Nguyen Tai Hoang, Nguyen Thi Tra My, Bui Lap Duy, Tran Van Hai Nam, Ha Thi Bich Hang, Le Thi Thuy Hang, Dinh Minh Hiep - Effect of rice bran oil on mycelial biomass production, biosynthesis and bioactivities of polysaccharides by <i>Ophiocordyceps sinensis</i> fungus.	74
11	Hoang Van Trung, Nguyen Thi Bich Ngoc, Nguyen Ngoc Tuan, Nguyen Tan Thanh, Tran Dinh Thang - Characterization of cytochalasins and steroids from the fruiting bodies of <i>Daldinia concentrica</i> in Viet Nam.	83
12	Nguyen Huu Tung, Tien Thi Luot, Nguyen Dang Linh, Cao Thi Mai Duyen - Entrainer selection for separation of azeotropic mixtures by distillation methods.	89
13	Ta Thi Lien, Nguyen Hoang Bao Tran, Nguyen Thi Thu Thuong, Nguyen Thi Hoa, Nguyen Thi Hoai Thu, Huynh Bui Linh Chi, Pham Nguyen Kim Tuyen, Nguyen Kim Phi Phung - Structural elucidation of five phenolic compounds from the rhizomes of <i>Curculigo annamitica</i> .	99
14	Pham Thi Nhat Trinh, Pham Ngoc An, Nguyen Trong Tuan, Nguyen Thi Le	106

- Thuy, Tong Thanh Danh, Le Tien Dung** - Research on phytochemical and alfa-glucosidase inhibitory activity of ethyl acetate fraction of *Ruellia tuberosa*.
- 15 **Nguyen Thi Hien, Hoang Dinh Hoa, Nguyen Quynh Hoa, Pham Thi Tram, Nguyen Thi Minh Tu** – Study on extraction of tannins from the *Garcinia mangostana* peel in Viet Nam. 113
- 16 **Nguyen Anh Hung, Dinh Thi Ha, Pham Quoc Long, Alla A. Kicha, Tran Thi Thu Thuy** - Steroidal diglycosides from the starfish *Anthenea sibogae*. 121
- 17 **Truong Ngoc Thao, Vu Nguyen To Nhu, Nguyen Thi Nguyen, Dong Thi Anh Dao** - Extraction of chondroitin sulfate from chicken keel cartilage by combined application of ultrasound treatment and alcalase hydrolysis. 127
- 18 **Nguyen Van Cong, Do Thi Hoa Vien, Dang Diem Hong** - Fatty acid profile and nutrition values of the microalga (*Thalassiosira pseudonana*) used in white shrimp culture. 138
- 19 **Hoang Duc Thuan, Nguyen Phi Hung, Vu Quoc Trung** - Methoxyflavones from *Orthosiphon stamineus* and their PTP1B inhibitory activities. 146
- 20 **Dinh Huy Son, Nguyen La Anh, Dang Thu Huong, Ly Ngoc Tram** - Effective method of polyphenol extraction from sesame cake using microbial fermentation by *Lactobacillus brevis* NCTH24. 153
- 21 **Le Thi Bach, Le Tien Dung, Nguyen Trong Tuan, Nguyen Quoc Chau Thanh, Joëlle Quetin-Leclercq, Bui Thi Buu Hue** - The protective effect of some extracts and isolated compounds from *euphorbia hirta* on pancreatic β -cells MIN6. 163
- 22 **Tran Quoc Hoa, Tran Thi Phuong Mai, Hoang Minh Nam, Nguyen Huu Hieu** - Extraction of polysaccharides from Lingzhi by ultrasonic-assisted enzymatic method. 171
- 23 **Nguyen Dang Binh Thanh, Nguyen Trung Dung, Ta Hong Duc** - Modeling of essential oil extraction process: application for orange, pomelo, and lemongrass. 182
- 24 **Truong Quoc Phong, Pham Thi Ngoc, Nguyen Dieu Huong, Nguyen Thi Ngoc Anh** - Efficient conjugation of aflatoxin B1 with bovine serum albumin for the development of aflatoxin B1 quick test. 190
- 25 **Le Xuan Duy, Tran Quoc Toan, Do Huu Nghi, Le Tat Thanh, Vu Dinh Hoang, Young Ho Kim, Nguyen Manh Cuong** – Triterpene acids from *Docynia indica* fruits and their cytotoxic activity. 199
- 26 **Cung Thi To Quynh and Vu Thu Trang** - Antimicrobial activities of Vietnamese holy basil (*Ocimum sanctum*) essential oil against food-borne bacteria and fungi. 205
- 27 **Vu Thi Thu Le, Do Tien Lam, Pham Minh Quan, Nguyen Thi Hong Van, Cam Thi Inh, Ta Thi Thu Thuy, Pham Thi Hong Minh** - Terpene compounds isolated from *Callicarpa macrophylla* in Viet Nam. 213
- 28 **Nguyen Tien Thanh, Le Thi Lan Chi, Do Thi Thu Ha, La Thi Quynh Nhu, Tran Thi Phuong Thao, Truong Quoc Phong, Khuat Huu Thanh** - Extraction of adenosine and cordycepin from spent solid medium of *Cordyceps militaris* culture. 221
- 29 **Nguyen Trung Dung, Ta Hong Duc, Nguyen Dang Binh Thanh** - Optimization of ginger oleoresin extraction from fresh ginger by using microwave-assisted energy. 209

- 30 **Mai Thi Thom, Tran Ngoc Toan, Tran Thi Thanh Huyen, Mai Van Chung** - 238
Change in content of glycosylated flavonoids in soybean (*Glycine max* cv. Namdan) leaves under aphid infestation.
- 31 **Do Xuan Hung, Luu Hang Nga, Tran Dinh Thang, Dang Ngoc Quang** - 246
Chemical constituents of the ethyl acetate fraction of the fruit bodies of *Phellinus gilvus*.
- 32 **Ngu Truong Nhan, Ninh The Son, To Dao Cuong, Nguyen Phuong Dai** 252
Nguyen, Pham Ngoc Khanh, Tran Thu Huong, Nguyen Manh Cuong -
Further study on chemical constituents from the heartwood of *Dalbergia tonkinensis* collected in Daklak province.
- 33 **Nguyen Thi Dieu Thuan, Nguyen Thi Thu Hien, Tran Minh Hao, Pham** 259
Van Huyen, Nguyen Huu Toan Phan - Flavonoids from the leaves of *Aralia hiiepiana*.
- 34 **Dinh Thi Thu Thuy, Tran Thi Tuyen, Nguyen Quyet Chien, Tran Thi Thu** 266
Thuy, Hoang Thi Bich, Tran Quoc Toan - Comparative study on volatile compounds of agarwood from Khanh Hoa province extracted by different methods.
- 35 **Tran Thi Minh, Do Thi Bien, Nguyen Thi Minh Thuong, Tran Thu Huong** - 273
Study on chemical constituents and biological activities of *Alpinia kwangsiensis* collected in Viet Nam.
- 36 **Ninh Thi Ngoc, Tran Thi Hong Hanh, Nguyen Xuan Cuong, Nguyen Van** 279
Thanh, Do Thi Thao, Nguyen Hoai Nam, Do Cong Thung, Phan Van Kiem, Chau Van Minh - Cholestane-type steroids from the octocoral *Verrucella corona*.

CHANGE IN CONTENT OF GLYCOSYLATED FLAVONOIDS IN SOYBEAN (*GLYCINE MAX* CV. NAMDAN) LEAVES UNDER APHID INFESTATION

Mai Thi Thom¹, Tran Ngoc Toan¹, Tran Thi Thanh Huyen², Mai Van Chung^{1,*}

¹ Vinh University, 182 Le Duan, Vinh city, Nghe An province, Vietnam

² Ha Noi University of Education, 136 Xuan Thuy, Cau Giay district, Hanoi, Vietnam

*Email: chungmv@vinhuni.edu.vn

Received: 12 July 2018; Accepted for publication: 1 October 2018

ABSTRACT

Changes in contents of glycosylated flavonoids in leaves of soybean (*Glycine max* (L.) Merr. cv. Namdan) in the flowering stage under infestation of cowpea aphid (*Aphis craccivora* Koch) were investigated. A strong decrease in the level of glycosylated flavonoids, e.g., genistein 7-O-glucoside, 2'OH genistein 7-O-glucoside, 2'OH genistein malonyl glucoside, after 48-96 hours post-infestation (hpi) of aphid was recorded. Levels of these compounds in aphid-infested leaves were remarkably lower than those in controls. Whereas, genistein and total flavonoids in the aphid-infested leaves were induced to higher contents than that observed in control. Activity of important enzymes regarding biosynthesis of flavonoids such as phenylalanine ammonia-lyase (PAL, EC. 4.3.1.5), chalcone synthase (CHS, EC 2.3.1.74) and chalcone isomerase (CHI, EC 5.5.1.6) in the aphid-infested leaves strongly increased during experimental time. An increase in activity of those enzymes was proportional with the induced decrease of the glycosylated flavonoids and an accumulated increase of genistein and total flavonoids. These results indicate that, changes in the glycosylated flavonoids occurred in Namdan soybean leaves infested by cowpea aphid may connect to a protective mechanism in this soybean cultivar in the flowering stage.

Keywords: Namdan soybean, cowpea aphid, glycosylated flavonoids, enzymes.

1. INTRODUCTION

Flavonoids have a variety of biological activities in plants, of which, the defensive functions to protect plants from different biotic and abiotic stresses were surely recorded [1]. The glycosylation of those antioxidants forming the glycosylated flavonoids impact several specific biochemical and physiological properties, which allows them to participate in plant defensive responses. The glycosylated flavonoids have been known to enable the plant defense mechanisms to better cope with biotic and abiotic stresses [2].

In our previous report [3], an accumulation of flavonoids was documented to participate in the defense responses of soybean (*Glycine max* (L.) Merr. cv. Namdan) to infestation of cowpea

aphid - *Aphis craccivora* Koch (Hemiptera: Aphididae). However, to date there has been lack of the available information of the insecticidal properties towards phloem-feeding insects of the glycosylated forms of flavonoids in this cultivar.

The aim of the present study was to examine whether genistein, a major component in soybean plants [4], and its glycosylation products such as genistein 7-O-glucoside, 2'OH genistein 7-O-glucoside and 2'OH genistein malonyl glucoside may be connected with responses of the Namdan soybean to effects of *A. craccivora* in the flowering stage. Simultaneously, post-infestation flavonoid contents were determined together with activity of enzymes regarding flavonoid metabolism such as phenylalanine ammonia-lyase (PAL), chalcone synthase (CHS) and chalcone isomerase (CHI).

2. MATERIALS AND METHODS

2.1. Materials

2.1.1. Plant

Plant used in this study is cultivar Namdan soybean (*Glycine max* (L.) Merr. cv. Namdan) that was cultured in Hoagland medium and placed in Plant physiology lab, Vinh University. The experimental conditions were such that: temperature of 23-25 °C, relative humidity of 70-75 %, light intensity of 110-130 $\mu\text{M photons.m}^{-2}.\text{s}^{-1}$ and light period of 14 light/10 dark hours. Soybean plants in the R3 stage (flowering stage) were used to establish experiments.

2.1.2. Aphid

Aphid species used in experiments is cowpea aphid (*Aphis craccivora* Koch), which is supported by Department of Applied Entomology (Institute of Ecology and Biological Resources, Vietnam Academy of Science and Technology).

2.2. Experiment

Each soybean plant in stage R3 (the flowering stage) was treated by 30 wingless adults of *A. craccivora*. This number of cowpea aphid was observed to cause the most serious damages on soybean plants [3]. Control was soybean plants without aphid infestation.

Control and aphid-infested variants were separately put in clear glass boxes (100 cm \times 50 cm \times 50 cm) covered by nylon gauze and placed in the growth chamber, in which, the cultured environmental factors such as temperature, relative humidity, light intensity and light period were strictly controlled.

Leaves of soybean were carefully collected after 0, 24, 48, 72 and 96 hours post-infestation (hpi) of cowpea aphid, then were weighed and frozen in nitrogen liquid for subsequent analyses.

2.3. Methods of analysis

2.3.1. Chemicals

All analytical chemicals were purchased from Singapore supplier of Sigma-Aldrich (USA).

2.3.2. Analysis of glycosylated flavonoids

Prior to LC profiling of glycosylated flavonoids, the frozen soybean leaves (0.5 g) were homogenized in 80 % methanol (20 mL g⁻¹FW) and sonicated for 3 min using a Bransonic 5510 bath sonicator. The suspension was filtered through a Büchner funnel and concentrated under vacuum at 40 °C. The samples were concentrated by solid phase extraction (SPE) on cartridges containing a cation exchanger and RP C18 silica gel (Alltech, Carnforth, England) [5].

Quantitative analyses were performed using an L7000 Merck Hitachi HPLC pump, equipped with an L7450 diode array detector (Darmstadt, Germany) and a Superspher 100 RP-18 column (250 mm × 2 mm; Merck). The amount of 125 mM p-hydroxybenzoic acid was added to each analysed sample as an internal standard.

Content of genistein, genistein 7-O-glucoside, 2'OH genistein 7-O-glucoside, and 2'OH genistein malonyl glucoside content was quantified by integrating UV chromatograms normalized at 259 nm to the peak of the standard. Their concentrations were expressed as milligram per gram fresh weight (mg.g⁻¹ fw) [6].

2.3.3. Analysis of total flavonoids

Content of total flavonoids in soybean leaves extract was determined by the aluminium chloride colorimetric method [7]. Total of 0.50 g frozen soybean leaves was extracted with 10 mL of 99.5 % ethanol under 200 rpm shaking for 24 hours. The filtrate was adjusted to 10 mL with 80 % ethanol and centrifuged at 10,000 × g for 10 min at 4 °C, the supernatant was collected and the precipitate was then extracted twice with 5 mL of 80 % ethanol.

A mixture of 200 µL extract and 150 µL of sodium nitrite (NaNO₂ 5 %, w/v), was incubated at room temperature for 6 min. Next, 150 µL of aluminium chloride hexahydrate (AlCl₃.6H₂O 10 %, w/v) was added and incubated for 6 min, then 1.000 µL NaOH (10 %, w/v) solution was added. For blank, distilled water was used to replace the extract. The absorbance was measured at λ = 510 nm in the UV-Vis CARY 60 spectrophotometer (Agilent, USA).

2.3.4. Enzymatic assays

Activity of phenylalanine ammonia-lyase (PAL, EC 4.3.1.5) was determined using the spectrophotometric method [8]. 0.50 g frozen soybean leaves was ground and extracted at 4 °C using 5 mL of 100 mM Tris-HCl buffer, pH 8.9, which contain 10 mM mercaptoethanol and 30 mg Polyclar AT. Mixture was centrifuged at 15,000 × g for 30 min at 4 °C. A volume of 1.5 mL of incubation mixture contained 80 mM borate buffer (pH 8.9), 30 mM L-phenylalanine and 0.5 mL enzymatic extract. The reaction was proceeded for 1 h at 30 °C and was halted by the addition of an equal volume of 2N HCl. The product of the reaction was assessed at 290 nm using an Agilent Cary 60 UV-Vis spectrophotometer.

Chalcone synthase (CHS, EC 2.3.1.74) was determined using the spectrophotometric method [9]. 0.5 g of soybean leaves was extracted in 5 mL of 100 mM borate buffer (pH 8.8) added 1 mM 2-mercaptoethanol at 4 °C. Then, the solution was added 0.1 g Dowex resin and centrifuged at 15,000 × g for 10 min. Supernatant was transferred to a tube, added 0.2 g Dowex resin, and incubated for 20 min. Mixture was centrifuged at 15,000 × g for 15 min. A volume of 2 mL mixture including 100 µL supernatant, 10 mM potassium cyanide, Tris-HCl buffer (pH 7.8), 10 mg chalcone was mixed with enzymatic extract. The absorbance was measured at 370 nm in Agilent Cary 60 UV-Vis spectrophotometer.

Activity of chalcone isomerase (CHI, EC 5.5.1.6) was assayed by spectrophotometry based on the isomerization of naringenin chalcone and isoliquiritigenin at 390 nm [10]. The reaction was carried out at 25 °C using 50 µM substrate in DMSO, 50 mM phosphate buffer (pH 7.6) containing 1 mg bovine serum albumin, and an appropriate amount of enzymatic extract. The reaction was initiated by adding the enzyme and control reactions were carried out without enzyme. The rate of spontaneous reaction was subtracted from the total rate.

2.3.5. Statistical analysis

All analyses were performed in at least three replicates in three independent experiments. Analysis of variance (ANOVA) was applied to verify whether means from independent experiments within a given experimental variant were significant with level of significance $\alpha = 0.05$. Data shown in the figures are means and standard errors (s.e.) for each variant.

3. RESULTS AND DISCUSSION

3.1. Content of glycosylated flavonoids in leaves of Namdan soybean

Among various glycosylated flavonoids, genistein and its derivatives such as genistein 7-O-glucoside, 2'OH genistein 7-O-glucoside and 2'OH genistein malonyl glucoside are the important endogenous compounds because they may enhance bioactivity in plant responses to biotic and abiotic stresses [1, 2, 6]. Their chemical structures are shown in Figure 1.




Figure 1. The structural scheme of genistein and its derivatives. Genistein ($R_1 = R_2 = H$); Genistein 7-O-glucoside ($R_1 = \text{glucose (glc)}$; $R_2 = H$); 2'OH Genistein 7-O-glucoside ($R_1 = \text{glc}$; $R_2 = OH$); 2'OH Genistein malonyl glucoside ($R_1 = (6''\text{-O-malonyl})\text{glc}$; $R_2 = OH$).

The change in content of those compounds in leaves of Namdan soybean under the different environmental conditions are shown in Figure 2.

After 24 hours post-infestation (hpi) of cowpea aphid, a drop in content of glycosylated isoflavonoids (genistein 7-O-glucoside, 2'OH genistein 7-O-glucoside, 2'OH genistein malonyl glucoside) was particularly pronounced since 24 hpi (Fig. 2a-c). ANOVA showed that differences in the level of glycosylated flavonoids between aphid-infested leaves were highly statistically significant. In the control leaves, a different trend was observed. The levels of glycosylated flavonoids changed a little and always higher, whereas, genistein mostly was lower than that in the infested variant during experimental time.




Figure 2. Content of glycosylated flavonoids (a-c) and genistein (d) in leaves of Namdan soybean infested by *A. craccivora* and control.

Interestingly, a significant decrease in contents of glycosylated flavonoids such as genistein 7-O-glucoside, 2'OH genistein 7-O-glucoside, 2'OH genistein malonyl glucoside, in the aphid infested leaves was probably a consequence of hydrolysis of the glycoside bonds in these molecules, leading to the release of genistein, a biologically active aglycone (Fig. 2d). Level of genistein in aphid infested leaves increased since 48 hpi as the glucosides strongly reduced.

Infestation of cowpea aphid can trigger oxidative burst in cells/tissues and cause serious damages to soybean plants [12]. Genistein of legume plants exhibits a wide range of biological activity, of which it play a significant role of antioxidant property, and thus counter the effects of oxidative stress. Therefore, metabolism of glycosylated flavonoids to form genistein may be connected with an enhancement the defense responses of *G. max* cv. Namdan to effects of *A. craccivora*.

3.2. Content of total flavonoids in leaves of Namdan soybean

In aphid-infested soybean leaves, level of total flavonoids were continuously increased and were always higher than that in the control during experiment (Fig. 3). The highest content of

flavonoids was $118.02 \mu\text{g QE.g}^{-1} \text{ dw}$ at 96 hpi, which was by 2.08- and 1.76-fold higher than at the beginning of experiment and in control plants, respectively. Whereas, content of flavonoids in control was minor changed during experimental time. The significant differences between flavonoids levels in the aphid-infested leaves and control were recorded from 72-96 hpi ($P < 0.05$). That result provided additional evidence about the defensive function of flavonoids in the defense mechanism of Namdan soybean against *A. craccivora* [3].




Figure 3. Content of total flavonoids in leaves of Namdan soybean infested by *A. craccivora* and control.

3.3. Activity of flavonoid-metabolic enzymes in leaves of Namdan soybean

During the experimental time, activity of phenylalanine ammonia-lyase (PAL, EC. 4.3.1.5), chalcone synthase (CHS, EC 2.3.1.74), and chalcone isomerase (CHI, EC 5.5.1.6) was observed to result in a strong increase in aphid-infested leaves of Namdan soybean, whereas, activity of those enzymes slightly changed in control plants (Fig. 4a). The highest activity of all enzymes was recorded at 96 hpi, which was from 2.08 (CHI)- and 2.51-fold (PAL) higher than that observed in control, respectively.

The biosynthesis pathway of flavonoids begins at precursors as phenylalanine catalyzed by PAL to form p-coumaroyl-CoA, which to yield chalcone (4',2',4',6'-tetrahydroxychalcone) catalyzed by CHS. The next step is isomerization of chalcone to flavanone by CHI. From this step onwards, the pathway branches to several different flavonoid classes including genistein and its glycosylate substances. An increase in activity of PAL, CHS, and CHI resulted in a strong increase in content of genistein (Fig. 2d) and total flavonoids (Fig. 3) in soybean leaves after aphid infestation, therefore, contributing to improve the defensive functions of those active substances in defense responses of *G. max* cv. Namdan to *A. craccivora*.




Figure 4. Activity of flavonoid-metabolic enzymes PAL (a), CHS (b) and CHI (c) in leaves of Namdan soybean infested by *A. craccivora* and control.

4. CONCLUSION

Infestation of *A. craccivora* induced the defense responses regarding flavonoids metabolism in leaves of soybean (*Glycine max* cv. Namdan) in the flowering stage. An increase in activity of important enzymes regarding biosynthesis of flavonoids such as PAL, CHS, and CHI was strongly connected with a strong decrease in the level of glycosylated flavonoids, e.g., genistein 7-O-glucoside, 2'OH genistein 7-O-glucoside, 2'OH genistein malonyl glucoside, and an accumulation of genistein and total flavonoids. Those changes in glycosylated flavonoids occurred in leaves infested by cowpea aphid were probably connected to the protection mechanism in Namdan soybean.

REFERENCES

1. Samanta A., Das G., Das S. K. - Role of flavonoids in plants, *Int. J. Pharm. Sci. Tech.* **6** (1) (2011) 12-35.
2. Ringli C., Bigler L., Kuhn B. M., Leiber R. M., Diet A., Santelia D., Frey B., Pollmann S., Klein M. - The modified flavonol glycosylation profile in the *Arabidopsis* roll mutants

- results in alterations in plant growth and cell shape formation, *The Plant cell* **20** (2008) 1470–1481.
3. Mai Van Chung, Doan Manh Dung, and Do Ngoc Dai - Accumulation of flavonoids in soybean under effects of Cyanobacteria crude extract and aphid infestation, *Vietnam Journal of Science and Technology* **54** (2C) (2016) 271-277.
 4. Dixon R. A., and Ferreira D. - Genistein, *Phytochemistry* **60** (3) (2002) 205-211.
 5. Stobiecki M., Wojtaszek P., and Gulewicz K. - Application of solid phase extraction for profiling of quinolizidine alkaloids and phenolic compounds in lupin tissue, *Phytochem. Anal.* **8** (1997) 153-158.
 6. Franski R., Bednarek P., Wojtaszek P., Stobiecki M. - Identification of flavonoid diglycosides in yellow lupin (*Lupinus luteus* L.) with mass spectrometric techniques, *J. Mass Spectrom.* **34** (1999) 486-495.
 7. Chang C. C., Yang M. H., Wen H. M., and Chern J. C. - Estimation of total flavonoid content in propolis by two complementary colorimetric methods, *J. Food Drug Anal.* **10** (2002) 178-182.
 8. Cahill D. M., and McComb J. A. - A comparison of changes in phenylalanine ammonialyase activity. Lignin and phenolic synthesis in the roots of *Eucalyptus callophylla* (field resistant) and *E. marginata* (susceptible) when infected with *Phytophthora cinnamoni*, *Physiol. Mol. Plant Pathol.* **40** (1992) 315–332.
 9. Ghasemzadeh A., Nasiri A., Jaafar H. Z. E., Baghdadi A., and Ahmad I. - Changes in phytochemical synthesis, chalcone synthase activity and pharmaceutical qualities of Sabah snake grass (*Clinacanthus nutans* L.) in relation to plant age, *Molecules* **19** (11) (2014) 17632-17648.
 10. Lister C. E., and Lancaster J. E. - Developmental changes in enzymes biosynthesis in the skins of red and of flavonoid green apple cultivars, *J. Sci. Food Agric.* **71** (1996) 313-330.
 11. Shimoda K., Kubota N., Hamada H., and Hamada H. - Synthesis of gentiooligosaccharides of genistein and glycitein and their radical scavenging and anti-allergic activity, *Molecules*, **16** (2011) 4740-4747.
 12. Tran Ngoc Toan, Tran Thi Thanh Huyen, Mai Van Chung - Antioxidative defense responses to aphid-induced oxidative stress in *Glycine max* (L.) Merr. cv. Namdan, *Vietnam Journal of Science and Technology* **54** (6) (2016) 719-728.